

Diagnosis and Symptoms of Narcolepsy From the Patient Perspective: Results From In-Depth Qualitative Interviews

Michael J. Doane¹; Meaghan O'Connor²; Miranda Lauher-Charest²; Laura Tesler Waldman²; and Wilbur P. Williams, III¹
¹Alkermes, Inc.; ²QualityMetric Incorporated, LLC

INTRODUCTION

- Narcolepsy is a rare, chronic neurological condition characterized by excessive daytime sleepiness (EDS) and sleep-onset rapid eye movement periods^{1,2}
- Narcolepsy is categorized into narcolepsy type 1 (NT1) and narcolepsy type 2 (NT2)²
 - Both types are characterized by EDS (including sleep attacks), sleep inertia, and sleep paralysis and/or hallucinations²
 - In addition, NT1 features cataplexy, which is a sudden, spontaneous, and temporary loss of muscle control triggered by strong emotional stimuli (e.g., fear, anger, laughter, or stress)^{2,3}
- Symptoms of narcolepsy often start in adolescence or early adulthood, but can occur at any time⁴
- Lack of symptom recognition by clinicians may lead to misdiagnosis, potentially delaying effective treatment and further exacerbating disease burden⁵
- Previous studies that identified symptoms of narcolepsy have largely relied on quantitative methods such as surveys, which may provide limited insight into the patient experience compared with qualitative methods

OBJECTIVE

- The objective of this study was to use qualitative research methods to characterize the patient experience of adults living with NT1 and NT2
- The data presented herein reflect the diagnosis journey and symptom burden of narcolepsy


METHODS

- This was a qualitative, cross-sectional, observational study
 - Using a semi-structured interview guide, interviewers asked open-ended questions to gather insights into people's experiences with NT1 or NT2
- Adult participants with NT1 or NT2 were recruited from panels of patients, physician referrals, and social media outlets
 - Snowball sampling, whereby participants could refer other participants to this study, was also used
- Interview transcripts were coded and thematically analyzed using inductive and deductive approaches
- This study was approved by an Institutional Review Board and all participants provided informed consent

RESULTS

PARTICIPANT DEMOGRAPHICS AND DISPOSITION


Figure 1. Demographics and Clinical Characteristics Among Participants



- A total of 22 adults with narcolepsy participated in this study (NT1 = 12; NT2 = 10)
- Overall, more than 70% of participants were female, and more than 60% of participants were White
- 83% of participants with NT1 and 50% of participants with NT2 were diagnosed with narcolepsy within the last 10 years

JOURNEY TO DIAGNOSIS


Figure 2. Time From Symptom Onset to Narcolepsy Diagnosis



- Symptom onset occurred most often in adolescence (n = 10, 45%) or early childhood (n = 6, 27%) among patients overall
- Initial symptoms typically included EDS (NT1 = 83%; NT2 = 80%), fatigue (NT1 = 42%; NT2 = 30%), oversleeping (NT1 = 33%; NT2 = 20%), sleep attacks (NT1 = 35%; NT2 = 50%), insomnia (NT1 = 35%; NT2 = 20%), and cataplexy (NT1 = 42%)
- 55% of patients overall (n = 12) reported that it took over 10 years for a diagnosis of NT1 or NT2 from symptom onset
- Half of participants (n = 11, 50%) described being misdiagnosed prior to their diagnosis of NT1 or NT2; misdiagnoses included depression (27%), sleep apnea (18%), and attention-deficit/hyperactivity disorder (9%)
- While seeking a diagnosis, participants reported visiting different types of healthcare professionals, including sleep specialists (NT1 = 7; NT2 = 4), neurologists (NT1 = 3; NT2 = 3), pulmonologists (NT1 = 2; NT2 = 2), psychiatrists (NT1 = 2; NT2 = 0), and primary care physicians (NT1 = 2; NT2 = 2)

FREQUENCY OF SYMPTOMS

Figure 3. Frequency of Symptoms Identified by Participants




- Almost all participants (n = 21; 95%) described experiencing EDS
 - Over three-fourths of participants (n = 18; 82%) reported sleep attacks, which tended to be intense, sudden, and uncontrollable
- 82% of participants (n = 18) reported fatigue, frequently describing it as a feeling of low or depleted energy that left them constantly "exhausted," "dragging," and "drained" during the day
- Nearly all participants (n = 21; 95%) described cognitive impairments, such as feeling "slow" or "foggy"
 - More than half of participants described lapses in short-term memory (NT1 = 9; NT2 = 5) and difficulties with focus and attention (NT1 = 5; NT2 = 7); nearly half described trouble thinking through and processing information (NT1 = 6; NT2 = 3)
- All participants with NT1 reported experiencing cataplexy, commonly triggered by strong emotions/reactions such as laughter (n = 6; 50%), anger (n = 5; 42%), stress (n = 5; 42%), and surprise (n = 3; 25%)
 - Almost all participants with NT1 (n = 11; 92%) described experiencing cataplexy within a specific body part, such as the legs, hands, head, neck, or face
 - One-third of NT1 participants (n = 4; 33%) described full-body cataplexy, often resulting in them collapsing to the ground
 - Frequency of cataplexy varied, ranging from a couple times a year to weekly or daily
- For those that could identify their most bothersome symptom, fatigue was the most frequently cited (NT1 = 5; NT2 = 4), followed by EDS (NT1 = 5; NT2 = 2), and cognitive impairment (NT1 = 2; NT2 = 1)
 - Those who selected fatigue described it as constant, severe, and debilitating

FROM THE PATIENT PERSPECTIVE

Figure 4. Patient-Reported Symptoms: In Their Own Words

The 4 most common symptoms were EDS, cognitive impairment, fatigue, and cataplexy (NT1 = 100%).



Study Limitations

- The rare nature of narcolepsy resulted in difficulty recruiting study participants
 - Despite this, a saturation analysis suggested saturation of concepts was reached for NT1 and NT2 participants and additional interviews would likely not have yielded new information
- Participants included in this study were mostly female, White, and under 40 years of age
 - It is unclear if results would have been different with a more diverse sample with respect to gender, race/ethnicity, and age
- Confirmation of diagnosis was encouraged but not required; confirmation was provided by 4 participants with NT1 and 8 participants with NT2

CONCLUSIONS

- Results of this study provide a comprehensive description of the patient experience of narcolepsy and highlight the often long and complicated journey these individuals experience while seeking a diagnosis for their varied symptoms
 - Most patients reported that they had been misdiagnosed since the onset of their narcolepsy symptoms, and it took the majority of patients over 10 years to receive a diagnosis of NT1 or NT2
- The rich descriptions provided by participants for each symptom have contributed to a deeper understanding of how each may be experienced in patients' own words
 - From the patient perspective, NT1 and NT2 are characterized by a constant desire to sleep during the day; challenges with focus, concentration, and memory; and overwhelming and debilitating fatigue
 - Additionally, NT1 includes cataplexy that ranges in frequency, location, and emotional triggers



REFERENCES

1. Ruoff C and Rye D. *Curr Med Res Opin.* 2016;32(10):1611-1622. 2. Sateia MJ. *Chest.* 2014;146(5):1387-1394. 3. Swick TJ. *Nat Sci Sleep.* 2015;7:159. 4. Dauvilliers Y, et al. *Neurology.* 2001;57(11):2029-2033. 5. Quaedackers L, et al. *Nat Sci Sleep.* 2021;13:1083-1096.

ACKNOWLEDGMENTS

The study was supported by Alkermes, Inc. Medical writing support was provided by Envision Pharma Group and was funded by Alkermes, Inc. This poster was developed in accordance with Good Publication Practice (GPP4) guidelines. Authors had full control of the content and made the final decision on all aspects of this poster.

DISCLOSURES

MJD and WPW are employees and stockholders of Alkermes, Inc. MOC, ML-C, and LTW are employees of QualityMetric Incorporated, LLC.

